

A comparative phytochemical investigation and antioxidant activity of *Quisqualis indica* L.

Hadeel M. Habeeb^{1*} 

¹Department of Biology, College of Science for Women, University of Baghdad, Iraq.

*Corresponding Author: Hadeel.makki@csw.uobaghdad.edu.iq

Article history:

Received: 17 December 2023

Accepted: 29 April 2024

Published: 30 June 2024

Keywords: *Quisqualis indica* L., Antioxidant, Flavonoids, HPLC, DPPH.

Abstract

Quisqualis indica L. is an herbaceous plant with many active ingredients that function as antioxidants and anticancer agents. Traditional medicine has long employed *Quisqualis indica* L. as an anti-inflammatory, anti-diarrheal, and anti-diabetic plant. The study aims to compare the antioxidant activity of leaves and flowers. By employing a particular reagent, a qualitative examination revealed the presence of alkaloids, terpenoids, tannins, flavonoids, and saponins. Flavonoids and phenolic compounds were isolated from leaves and flowers such as hydroxybenzoic acid, gallic acid, catechin, Rutin, kaempferol, and sinapic acid. According to the HPLC analysis, flowers contain a higher percentage of flavonoids and phenolic compounds than leaves. For the ascorbic acid, the antioxidant activity shows that the plant has significant inhibition activity against free radicals in both the leaves and the flowers, and the flowers are more active than the leaves. Also, the antioxidant activity increases with increasing concentration from 25 to 400 mg mL⁻¹; the IC₅₀ values for the flowers were 100 µg mL⁻¹ and 200 µg mL⁻¹ for the leaves.

<https://doi.org/10.52951/dasj.24160112>

This article is open-access under the CC BY 4.0 license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

In ancient times, herbal medicines were the primary source of treatment for many diseases (Ssenku *et al.*, 2022). Even today, in many places, they are still used for healthcare. Therefore, herbal medicine can be considered a traditional system of medicine that is used in medical practices since antiquity (Yuan *et al.*, 2016). formerly known as *Holarrhena floribunda*, displays a wide range of pharmacological activities such as anti-inflammatory, antipyretic, immunomodulatory, anti-staphylococcal, anthelmintic, and antiseptic activities. Plant components have distinct energy primary that cause the activities, *Quisqualis indica* is a combretaceae perennial vine (Riaz *et al.*, 2023).

Quisqualis indica plants are distinguished by oval leaves with parallel veins (Sahu *et al.*, 2012). The blooms range in color from pale

pink to deep pink or crimson, according to Kulshreshtha *et al.* (2018). The plant's leaves and flowers include alkaloids, flavonoids, saponins, and tannins (Barik *et al.*, 2020). Leaves have more terpenoids than flowers, which have more phenols. Thu *et al.* (2022) found that oils are in the flowers and triterpenes, sterols, and aromatic chemicals in the leaves. (Mohammed and Habeeb, 2022).

Abd El Rahman *et al.* (2008); Revathi and Radh (2016); and Nemade and Aher (2023) have proved that plants improve health, with many benefits like Acetylcholinesterase inhibition, cholesterol reduction, fever reduction, worm prevention, antioxidants, inflammation reduction, infection prevention, and cancer prevention. The presence of stress in cells can be triggered by many factors like smoking, pollution, pesticides, or internal reactive oxygen species (ROS) such as

superoxide anion, singlet oxygen and hydrogen peroxide. Stress can lead to damage in acids, proteins and enzymes altering their integrity within living organisms as studied by González Palma *et al.* (2016). The body's antioxidant defense mechanisms work to minimize and stabilize these factors. Current research is focusing on the use of tannins and flavonoids, in supplements to combat free radical damage effectively and alleviate ROS effects according to Saeed *et al.* (2012).

The main challenges are to extract and isolate specific active groups from medicinal plants without removing other active chemicals (Ghenabzia *et al.*, 2023), and then assess the biochemical behavior and bioavailability, including toxicity (Sasidharan *et al.*, 2011) and *in vivo*. This paper aims to extract and investigate active compounds, then isolate flavonoids and phenolic compounds, conduct qualitative flavanol and phenolic compounds analysis, and examine their antioxidant activity.

Materials and Methods

Plant collection and extraction

The plant leaves and flowers were collected in November from Baghdad nurseries, and then washed. The leaves were dried, but the flowers were used freshly, the dry leaf was powdered and the flowers were extracted with a (1:10) ratio (plant: solvent) in a soxhlet with petroleum ether for 4 hours at 60–80 °C separately, then the plant material was extracted with methanol at 60–80 °C for 6–8 hours, then filtered (Ali *et al.*, 2022).

Qualitative analysis of an active compound

The active components investigated were detected according to Ali *et al.* (2022). The 1g of crud extract (leaves and flowers) was dissolved in 10 ml of methanol. 1 ml of this extract was directed to the detection of active compound presence, alkaloids, terpenoids,

tannins, flavonoids, and saponins using specific two different reagents for each compound, as follows:

1. Alkaloids: Mayer and Wagner reagents are used for the detection process.
2. Terpenoids: Two reagents, chloroform, and H₂SO₄, as well as an aldehyde reagent, were utilized to detect the terpenoids.
3. Tannins: by adding FeCl₃ reagent and lead acetate reagent.
4. Magnesium crystals with 1% HCl and saponins with H₂SO₄ reagent.
5. Saponins reagents for foam, HgCl₂, and other substances.

Phenolic compound and Flavonoids purification

The leaves and flowers methanolic extract were concentrated, then sequentially distilled water and ethyl acetate was added for each flower and leaf, and then shaken. After that, the organic layer was collected and dried. The presence of phenolic compounds and flavonoids was detected (Wagner and Bladt, 2009).

Quantitative analysis of phenolic compound flavonoids

The purified phenolic compound flavonoids were quantitatively determined by high-performance liquid chromatography (HPLC), which was done using the Sykamn Germany HPLC system, with C18-ODS column dimensions (250 mm * 4.6 mm, 5µm). 100 µm of samples were injected, and the sample was developed using acetonitrile and 0.01% trifluoroacetic acid at 1 mL min⁻¹ in the mobile phase at a 1 mL min⁻¹ flow rate. The developing program followed the sequence concentration (10% from 0–5 min; 25% from 5-7 min; 40% from 7–13 min) of acetonitrile, respectively, returning to the first concentration condition. Then a UV-visible detector at 278 nm was used

to detect the phenolic compounds, flavonoid concentrations calculated by the following equation (Ngamsuk *et al.*, 2019).

The concentration of flavonoids = [area of specimen/area of standard * concentration of standard * dilution factor].

Antioxidant activity

Leaf and flower phenolic compound flavonoids antioxidant abilities were examined through scavenging 1,1'-diphenyl-2-picrylhydrazyl (DPPH) according to Marwah *et al.* (2007) 2 mL of stock solution 100 mM DPPH in methanol was mixed with 2 ml of leaf or flower flavonoids to make the reaction medium. Ascorbic acid was used as a standard. After mixing, the mixture was incubated at 35 °C in the dark for 20 min. at 512 nm. The absorbance was then recorded, and the treatment was triple-repeated. The IC₅₀ (concentration of sample required to scavenge 50% of DPPH radicals) The DPPH scavenging ability is calculated based on the following equation:

$$\% \text{ DPPH radical scavenging activity} = [1 - (A_{\text{sample}}/A_{\text{control}}) * 100]$$

A control = absorbencies of control

A sample = absorbencies of sample

Statistical Analysis

The SAS program was previously used to determine different factors in the study of parameters. LSD test and ANOVA were used to significantly compare between means in this study.

Results and Discussion

Qualitative analysis of an active compound

The active compound test of *Quisqualis indica* leaf and flower crud extract shows the presence of alkaloids, terpenoids, tannins, flavonoids, and saponins in Table 1, this result goes with Barik *et al.* (2020) of presence of alkaloids, tannins, flavonoids, and saponins in leaf and flower extracts, while the terpenoids are found in leaves and absent in flowers, but Shah *et al.* (2019) shows the absence of alkaloids and tannins in *Quisqualis* leaves and flowers. The contrast in the presence or absence of some active compounds in the same genus may be related to the environment and where the plant is cultivated (Pant *et al.*, 2021).

Table 1. Phytochemical screening of active ingredients in *Quisqualis indica* leaf and flower crud extract

Active compounds	leave		flower	
	Reagent A	Reagent B	Reagent A	Reagent B
Alkaloids	+ white precipitate	+ brown precipitate	+ white precipitate	+ brown precipitate
Terpenoids	+ Brown-redush	+ brown precipitate	+ Brown-redush	+ brown precipitate
Tannins	+ green- blue	+ yellow precipitate	+ green- blue	+ yellow precipitate
Flavonoids	+ red- orange	+ red	+ red- orange	+ red
Saponins	+ foam	+white precipitate	+ foam	+white precipitate

+ Presence of active compound

Quantitative analysis of phenolic compound and flavonoids

The results of phenolic compound and flavonoids quantitative analysis after flavonoids

and phenolic compound isolation from crude extract, which is shown in (Table 2), indicate the presence of sex compounds in the purified extract, which show that the extract of leaves and flowers contains hydroxybenzoic acid, gallic acid, catechine, rutin, kaempferol, and

sinapic acid figure (1 and 2); Tables 3, 4 and 5, which refer to the retention time of each stander, leaves, and flowers. Table 3 shows that hydroxybenzoic acid, the same molecule, has a retention period of 2.51 for standard hydroxybenzoic acid and 2.58, and 2.55 for leaves and flowers, respectively. Stander gallic acid was 3.02, in addition, it was 3.05 and 3.01 for leaves and flowers, it was 3.05 and 3.01; in the stander category retention time, the leaves were 5.95, the flowers were 5.96, and the catechine stander was 5.90. The retention times for rutin, kaempferol, and sinapic acid are as follows: 6.52 for rutin, 7.85 and 7.81 for leaves and flowers, 9.35 for kaempferol, 9.32 and 9.35 for leaves and flowers, and 11.25 for sinapic acid stander and 11.26 and 11.27 for leaves and flowers, respectively. The approximate

retention times of the flowers, leaves, and standers are identical, suggesting that the isolated compounds are similar to the standers. The result shows that all phenolic compounds and flavonoids were higher in the flowers than the leaves, while gallic acid has the highest compound concentration and sinapic acid has the lowest amount among the active compounds in each flower and leaf. This result differs from Rastogi *et al.* (2019), which elucidate that the quantity of leaf flavonoids is higher than flower flavonoids and agrees with them in the presence of some active compounds (gallic acid and rutin), also the result is compatible with Bairagi *et al.* (2012), and Chaudhary *et al.* (2021) and in the presentation of some types of flavonoids in Iraqi plants.

Table 2. Type and Quantity of phenolic compound and flavonoids isolated from *Quisqualis indica* by HPLC chromatography

NO	Name	Leaves (ppm)	Flowers (ppm)
1	Hydroxybenzoic acid	12.8	15.1
2	gallic acid	33.9	42.6
3	catechine	22.5	29.8
4	Rutin	21.6	30.6
5	kaempferol	24.9	29.7
6	Sinapic acid	9.6	12.9

Table 3. The retention time and peak area of HPLC Chromatography of standers phenolic compound and flavonoids



No	Compounds	Retention time	Peak area
1	hydroxybenzoic acid	2.51	1421.05
2	gallic acid	3.02	1521.44
3	catechine	5.90	1244.79
4	Rutin	6.52	952.11
5	kaempferol	9.35	1877.48
6	Sinapic acid	11.25	952.14

Table 4. The retention time and peak area of HPLC Chromatography of *Quisqualis indica* Leaves

No	Compounds	Retention time	Peak area
1	hydroxybenzoic acid	2.58	2564.25
2	gallic acid	3.05	3652.14
3	catechine	5.95	1569.89
4	Rutin	7.85	2698.59
5	kaempferol	9.32	3655.49
6	Sinapic acid	11.26	6521.47

Table 5. The retention time and peak area of HPLC Chromatography of *Quisqualis indica* Flowers

No	Compounds	Retention time	Peak area
1	hydroxybenzoic acid	2.55	3652.89
2	gallic acid	3.01	5412.49
3	catechine	5.96	2145.88
4	Rutin	7.81	4875.80
5	kaempferol	9.35	5991.41
6	Sinapic acid	11.27	8562.08

**Figure 1. HPLC chromatography of leaves phenolic compound and flavonoids****Figure 2. HPLC chromatography of flowers phenolic compound and flavonoids**

Antioxidant activity assay

The results in Table 6 elucidate the antioxidant activity of leaves and flowers phenolic compound and flavonoids isolated from *Quisqualis indica* using the DPPH method and compares them with ascorbic acid as a moisturizer, which indicates the *in vitro* ability

of *Quisqualis indica* phenolic compound and flavonoids to scavenge the free radical factors released in solution. The results showed that scavenger activity increased with phenolic compound and flavonoids concentration increased for leaves and flowers and exceeded flower flavonoids on leaf flavonoids, and the

higher inhibition rate was 76.62 and 73.18 in concentration 400 $\mu\text{g/mL}$, while the lowest inhibition rate was 36.95 and 28.70 in 25 $\mu\text{g/mL}$ for flowers and leaves, respectively, while the IC_{50} was 100 $\mu\text{g/mL}$ for flowers (57.25) and 200 $\mu\text{g/mL}$ for leaves and ascorbic acid (66.08 and 54.80) respectively. The results presented significantly differ ($p < 0.001$) between all concentrations used in the experiment. The antioxidant activity of the *Quisqualis indica* plant agrees with Shah *et al.* (2019), which elucidate the alcoholic extract of *Quisqualis*

indica leaves and flowers. These antioxidant effects of leaf and flower phenolic compound and flavonoids due to the high flavonoid concentration, which has a strong antioxidant effect and ability to scavenge free radical factors because the secondary metabolites like phenols and flavonoids neutralize, absorb, and scotch O_3 and other free radicals due to their redox activity, structure of the conjugated ring, and presence of a carboxyl group (Olugbami *et al.*, 2014).

Table 6. Scavenging activity of flowers and leaves compound

Concentration $\mu\text{g} \cdot \text{mL}^{-1}$	Mean scavenging activity \pm SE		
	Ascorbic acid	Flowers	Leaves
25	20.98 \pm 0.84 e	36.95 \pm 1.62 d	28.704 \pm 1.27 c
50	31.48 \pm 1.78 d	43.94 \pm 2.36 d	41.165 \pm 2.08 b
100	41.04 \pm 2.37 c	57.25 \pm 2.89 c	48.30 \pm 2.62 b
200	54.80 \pm 2.95 b	68.01 \pm 3.55 b	66.088 \pm 3.27 a
400	63.503 \pm 3.02 a	76.62 \pm 3.82 a	73.18 \pm 3.08 a
LSD value	7.0216 **	8.441 **	7.594 **
P-value	0.0001	0.0001	0.0001

Means having the different letters in the same column differed significantly, ** ($P \leq 0.01$).

Conclusions

Active chemicals extracted from *Quisqualis indica* L. and other plants are being recognized as a new strategy for preventing and treating numerous human ailments. This research compares the chemical presence of alkaloids, terpenoids, tannins, flavonoids, and saponins in flowers and leaves of the same plant. High-performance liquid chromatography (HPLC) was used for quantitative analysis of phenolic compounds and flavonoids. The HPLC analysis revealed that the flowers contain a larger percentage of phenolic compounds and flavonoids compared to the leaves. The leaves and flowers exhibited considerable antioxidant activity against free radicals when compared to ascorbic acid. The antioxidant activity increased as the concentration increased, with flowers having an IC_{50} of 100 $\mu\text{g mL}^{-1}$ and leaves

having an IC_{50} of 200 $\mu\text{g mL}^{-1}$. This occurrence has led to a recent increase in demand for it.

Conflict of Interest

There are no disclosed conflicts of interest for authors.

Acknowledgements

The author would like to thank the College of Science for Women, University of Baghdad for their facilities in using the biology laboratories.

References

- Abd El-Rahman, A. A., Abd El-Aleem, I. M., Refahy, L. A., and El-Shazly, M. A. (2016). Total phenolic content, cytotoxic and antioxidant activities of *Quisqualis indica* (Linn.) growing in Egypt. *Der Pharmac Chemica*, 8(3), 53-9.

- Ali, Z. A. A., Habeeb, H. M., and Jazaa, L. A. (2022). Morphological, Anatomical and Chemical Study of an Exotic Plant *Jatropha integerrima* JACQ. 1763 (Euphorbiaceae) in Iraq. *Bulletin of the Iraq Natural History Museum*, 17(1), 129-140.
<https://doi.org/10.26842/binhm.7.2022.17.1.0129>
- Anjali, P., Vivek, J., Jigar, P., Keyuri, B., Vaibhavi, S., Jayesh, D., and Devang, P. (2023). Establishment of quality parameters of *Quisqualis indica* leaves through sophisticated analytical techniques. *Bulletin of the Chemical Society of Ethiopia*, 37(2), 289-298.
<https://doi.org/10.4314/bcse.v37i2.4>
- Bairagi, V. A., Shinde, P. R., Senthikumar, K. L., and Sandu, N. (2012). Isolation and characterizations of phytoconstituents from *Quisqualis indica* Linn. (Combretaceae). *Research Journal of Pharmacognosy and Phytochemistry*, 4(4), 229-233.
- Barik, B. S., Das, S., and Hussain, T. (2020). Pharmacognostic properties of *Quisqualis indica* Linn: against human pathogenic microorganisms: an insight review. *European Journal of Medicinal Plants*, 31(20), 87-103.
<https://doi.org/10.9734/ejmp/2020/v31i2030369>
- Chaudhary, D., Srivastava, R., and Nagar, H. (2021). Anti-allergic Assessment of Ethanol Extractives of *Quisqualis Indica* Linn. *Current Bioactive Compounds*, 17(7), 48-57.
<https://doi.org/10.2174/1573407216999201124222935>
- Ghenabzia, I., Hemmami, H., Ilham, B. E. N., Zeghoud, S., Seghir, B. B., and Hammoudi, R. (2023). Different methods of extraction of bioactive compounds and their effect on biological activity: A review. *International Journal of Secondary Metabolite*, 10(4), 469-494.
<https://doi.org/10.21448/ijsm.1225936>
- González-Palma, I., Escalona-Buendía, H. B., Ponce-Alquicira, E., Téllez-Téllez, M., Gupta, V. K., Díaz-Godínez, G., and Soriano-Santos, J. (2016). Evaluation of the antioxidant activity of aqueous and methanol extracts of *Pleurotus ostreatus* in different growth stages. *Frontiers in microbiology*, 7, 1099.
<https://doi.org/10.3389/fmicb.2016.01099>
- Jahan, F. N., Rahman, M. S., Hossain, M., and Rashid, M. A. (2008). Antimicrobial activity and toxicity of *Quisqualis indica*. *Advances in Traditional Medicine*, 8(1), 53-58.
<https://doi.org/10.3742/OPEM.2008.8.1.053>
- Kulshreshtha, M., Shukla, K. S., Tiwari, G. A., Singh, M. P., and Singh, A. (2018). Pharmacognostical, phytochemical and pharmacological aspects of *Quisqualis indica*: An update. *Journal of Nature and Science of Medicine*, 1(2), 41-47.
- Marwah, R. G., Fatope, M. O., Al Mahrooqi, R., Varma, G. B., Al Abadi, H., and Al-Burtamani, S. K. S. (2007). Antioxidant capacity of some edible and wound healing plants in Oman. *Food chemistry*, 101(2), 465-470.
<http://dx.doi.org/10.1016/j.foodchem.2006.02.001>
- Mohammed, S., and Habeeb, H. M. (2022). Phytochemical and cytotoxic study in *Quisqualis indica* volatile oil cultivation in Iraq. *International Journal of Health Sciences*, 6(S4), 8572– 8584.
<https://doi.org/10.53730/ijhs.v6nS4.10618>
- Nemade, C., and Aher, A. (2023). *Quisqualis* Estimation of Phytoconstituents and Anti-Histaminic Activity of *Quisqualis indica* Linn. *International Journal of Pharmaceutical Sciences and Research*, 14(5), 2429-2433.
[http://dx.doi.org/10.13040/IJPSR.0975-8232.14\(5\).2429-33](http://dx.doi.org/10.13040/IJPSR.0975-8232.14(5).2429-33)
- Ngamsuk, S., Huang, T. C., and Hsu, J. L. (2019). Determination of phenolic compounds, procyanidins, and antioxidant

- activity in processed *Coffea arabica* L. leaves. *Foods*, 8(9), 389.
<https://doi.org/10.3390/foods8090389>
- Olugbami, J. O., Gbadegesin, M. A., and Odunola, O. A. (2014). In vitro evaluation of the antioxidant potential, phenolic and flavonoid contents of the stem bark ethanol extract of *Anogeissus leiocarpus*. *African journal of medicine and medical sciences*, 43(1), 101-109.
<https://pubmed.ncbi.nlm.nih.gov/26681826/>
- Pant, P., Pandey, S., and Dall'Acqua, S. (2021). The influence of environmental conditions on secondary metabolites in medicinal plants: A literature review. *Chemistry and Biodiversity*, 18(11), e2100345.
<https://doi.org/10.1002/cbdv.202100345>
- Rastogi, S., Pandey, M. M., Rawat, A. K. S., Kushwaha, V., and Murthy, P. K. (2019). In vitro antifilarial activity, antioxidant potential and phenolic constituents of *Quisqualis indica* L. *Indian Journal of Traditional Knowledge*, 18(4), 648-654.
<http://op.niscair.res.in/index.php/IJTK/article/view/28936>
- Revathi, D., and Radha, K. (2018). Phytochemical screening of *Quisqualis indica* Linn. *Int. J. Innov. Res. Multidiscip. Field*, 4(8), 154-156.
- Riaz, M., Khalid, R., Afzal, M., Anjum, F., Fatima, H., Zia, S., and Aslam, M. A. (2023). Phytobioactive compounds as therapeutic agents for human diseases: A review. *Food Science and Nutrition*, 11(6), 2500-2529.
<https://doi.org/10.1002/fsn3.3308>
- Saeed, N., Khan, M. R., and Shabbir, M. (2012). Antioxidant activity, total phenolic and total flavonoid contents of whole plant extracts *Torilis leptophylla* L. *BMC complementary and alternative medicine*, 12(221), 1-12.
<https://doi.org/10.1186/1472-6882-12-221>
- Sahu, J., Patel, P. K., and Dubey, B. (2012). *Quisqualis indica* Linn: A review of its medicinal properties. *International Journal of Pharmaceutical and Phytopharmacological Research*, 1(5), 313-321.
- SAS. (2018). *Statistical Analysis System, User's Guide. Statistical. Version 9.6th. ed.* SAS. Inst. Inc. Cary. N.C. USA.
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K. M., and Latha, L. Y. (2011). Extraction, isolation and characterization of bioactive compounds from plants' extracts. *African journal of traditional, complementary and alternative medicines*, 8(1), 1-10.
<https://doi.org/10.4314/ajtcam.v8i1.60483>
- Shah, A., Saleem, S., and Farid, S. (2019). Antioxidant activity of an ethnobotanically important plant *Quisqualis indica* Linn. *Pakistan Journal of Pharmaceutical Sciences*, 32(1), 95-102.
- Ssenku, J. E., Okurut, S. A., Namuli, A., Kudamba, A., Tugume, P., Matovu, P., and Walusansa, A. (2022). Medicinal plant use, conservation, and the associated traditional knowledge in rural communities in Eastern Uganda. *Tropical Medicine and Health*, 50, 39, 1-10. <https://doi.org/10.1186/s41182-022-00428-1>
- Thu, N. T. H., Hoa, P. T. H., Dat, N. T., and Tuyen, P. N. K. (2022). Triterpenoids, steroid, and aromatic compounds from *Combretum indicum* leaves. *Vietnam Journal of Chemistry*, 60(5), 629-635.
<https://doi.org/10.1002/vjch.202100211>
- Wagner, H. and Bladt, S. (2009). *Plant Drug Analysis. Second edition.* Springer Dordrecht Heidelberg. London. New York.
- Yuan, H., Ma, Q., Ye, L., and Piao, G. (2016). The traditional medicine and modern medicine from natural products. *Molecules*, 21(5), 559.
<https://doi.org/10.3390/molecules21050559>