

# Effect of Dietary Intake of Lutein Rich Foods on the Retina and its Associated with Retinopathy Among Diabetic Patients

Hiba Raad Saeed <sup>1</sup>, Ahmed Ghazi Dadoosh <sup>2</sup>, Besmah Mohamad Ali <sup>3</sup>, Khaled Awad Mohamed Elbassiouny <sup>4</sup>

<sup>1</sup> Division of communicable disease control, Public health department, Baghdad Al-Rusafa Health Directorate, Baghdad, Iraq.

<sup>2</sup> Ophthalmology department, Imamein Kadhimein medical city, Baghdad Al-Karkh Health Directorate, Baghdad, Iraq.

<sup>3</sup> Public health department, Ghazi Al-Hariri Surgical Hospital, Baghdad, Iraq.

<sup>4</sup> Ophthalmology consultant, Lecturer of Ophthalmology, Delta University, New-Mansoura, Egypt.

## Abstract

**Background:** Diabetic retinopathy (DR) is a common micro-vascular consequence of diabetes mellitus that affects the eyes. If untreated, DR cause damage to the retinal blood vessels and result in blindness or visual loss. Hyperglycemia over long period of time results in increases inflammatory oxidative stress and protein kinase C pathways that eventually retinal capillary endothelial damage and pericyte loss occurred. Lutein is a carotenoid with anti-inflammatory and antioxidant properties.

**Objective:** To determine the protection roles of dietary intake of lutein rich foods on the retina and it's associated with retinopathy among diabetic patients.

**Patients and Methods:** A cross-sectional study was conducted in the Ophthalmology Department of Imamein Kadhimein Medical City, Baghdad, Iraq, from July to November 2023, involving a total of 100 diabetic patients. Questionnaire list was used which consist of sociodemographic information, socioeconomic status, dietary sources of lutein, anthropometric measures and ophthalmic examination and investigations which include optical coherence tomography for macular assessment and HbA1c assessment.

**Results:** The current study showed that 64% of the diabetic patients had retinopathy and 36% had normal retina. The study showed that 91.7% of diabetic patients with normal retina were consume 3-7 serving/week tomato, 75% were consume 7-35 serving/week egg, 75% were consume 1-7 serving/week zucchini, 72% were consume 1-3 serving/week spinach, 63.9% were consume 3-7 serving/week green pepper, 63.9% were consume 3-7 serving/week basil, 61% were consume 3-7 serving/week parsley, 58% were consume 3-7 serving/week leek, 52.8% were consume 3-7 serving/week lettuce, 41.7% were consume 1-2 serving/week nut and 30.6% were consume 1-2 serving/week pistachio with significant p-value.

**Conclusion:** Patients with diabetes can avoid retinal damage by consuming the recommended weekly servings of lutein-rich foods especially tomato, eggs, zucchini, spinach, green pepper, basil, parsley, leek, lettuce, nut and pistachio.

**Keywords:** Diabetic retinopathy, lutein, visual impairment.

**Correspondence Address:** Hiba Raad Saeed

**Email:** [hebars81@yahoo.com](mailto:hebars81@yahoo.com)

**Copyright:** ©Authors, 2024. College of Medicine, University of Diyala. This is an open access article under the [CC BY 4.0](http://creativecommons.org/licenses/by/4.0/) license (<http://creativecommons.org/licenses/by/4.0/>)

**Website:**

<https://djm.uodiyala.edu.iq/index.php/djm>

**Received:** 17 July 2024

**Accepted:** 03 December 2024

**Published:** 25 December 2024

## Introduction

The eye is an important organ that needs special care (1). Diabetes and its consequences are growing more prevalent worldwide (2). By 2030, it is anticipated that over 191 million individuals globally will suffer from diabetic retinopathy, with over 55 million of them experiencing visual impairment (2). There are 1.4 million type 2 diabetics in Iraq, where the disease affects 8.5% to 13.9% of the population (3). In Iraq, the prevalence of diabetic retinopathy ranged from 28.1% to 32.8%, with proliferative alterations observed in 11.27% of cases (4). Globally, 22.27% of people had DR (5). Diabetic retinopathy is a microvascular disorder that destroys retinal blood vessels and can lead to blindness or visual loss if untreated (6 – 8). Hyperglycemia over long period of time results in increased inflammatory oxidative stress and protein kinase C pathways that eventually retinal capillary endothelial damage and pericyte loss were occurred. Retinal capillary changes causing capillary occlusion, retinal non perfusion and endothelial barrier decompensation which leads to serum leakage and retinal edema (macular edema). These changes occur in both superficial and deep retinal capillary vessels and worsen in more severe diabetic retinopathy. Retinal neovascularization may develop due to increased level of intravascular endothelial growth factor (VEGF) from ischemic retinal tissue (7, 8). Diabetic retinopathy (DR) is a leading cause of vision loss globally, ranking sixth in terms of preventable blindness and fifth in terms of moderate-to-severe visual impairment between 1990 and 2020 (9). A third of individuals with diabetes mellitus have vision-

threatening retinopathy, which is characterized by either proliferative or severe non-proliferative DR or the presence of diabetic macular edema (7, 8, 10). Approximately one in three individuals with diabetes mellitus have DR (11). DR is estimated to be the most frequent cause of new cases of blindness among adults 20-74 years of age (12). Studies have demonstrated that daily ingestion of a multi-component formula combining antioxidants and xanthophyll pigments improves perimetry, color discrimination, macular pigment optical density, and contrast sensitivity in diabetic individuals with and without retinopathy (13). Lutein and zeaxanthin are carotenoids that have anti-inflammatory, antioxidant (14-16) and neuroprotective effects (17). Unique carotenoids that are concentrated in the human macula (center of retina) are: lutein, zeaxanthin, and meso-zeaxanthin (16-19). Lutein and zeaxanthin are obtained from dietary sources such as green leafy vegetables and orange and yellow fruits and vegetables, while meso-zeaxanthin is rarely found in diet and is believed to be formed at the macula by metabolic transformations of ingested carotenoids (17, 19). Several studies have proved that lutein and zeaxanthin are an essential element for eye health (1, 14 – 21). Elevated levels of the food-dependent plasma carotenoids lutein and zeaxanthin are thought to offer protection against diabetic retinopathy (13 – 15). Consuming lutein and zeaxanthin has been shown to help diabetic patients with non-proliferative retinopathy, macular edema and improve their contrast sensitivity and visual acuity (13 – 15). Lutein and zeaxanthin may have protective effect for macula in

patients with DR because of their biochemical structure and function that neutralize reactive oxygen species and prevent oxidative damage to the retina (biological antioxidants), neuroprotective and anti-inflammatory function in the retina, its position in the center of retina (macula), and its ability to absorb oxidative blue light (2, 14 – 17). Carotenoids that are present in large quantities in egg yolks, orange and yellow fruits, and dark green vegetables (Leek, Parsley, Lettuce, green pepper), tomato and nut are lutein and zeaxanthin (18, 20-22). It is also one of just two carotenoids that are present in the human eye, where it serves to shield the retina from damaging light and oxidation (18). Dietary factors have a significant impact on DR risk modification, as evidenced by the protective effects of a Mediterranean diet, high consumption of fruits, vegetables, and fish, and low calorie intake (22 – 24). Depending on a fact that lutein has antioxidant, anti-inflammatory and neuroprotective effects. Hence, the objective of this study was to determine the protection roles of dietary intake of lutein rich foods on the retina and it's associated with retinopathy among diabetic patients.

### Patients and Methods

The Department of Ophthalmology of Imamein Kadhimein Medical City, Baghdad, Iraq was the site for data collection of the current cross-sectional study, for five months duration from the 1<sup>st</sup> of July to 1<sup>st</sup> of December 2023. One-hundred diabetic patients were collected from outpatient in ophthalmology department of this hospital; Patients were only included if they met the inclusion criteria which are all diabetic patients aged between 35- 90 years.

Questionnaire list was used in current study and consist of sociodemographic information, socioeconomic status, and dietary sources of lutein, anthropometric measures and ophthalmic examination.

**1. Sociodemographic information** consist of age, sex, education, occupation and marital status.

**2. Socioeconomic status** was determined based on standard equation: Education + Occupation + house ownership  $\times 0.5$  + car ownership  $\times 0.1$  (25).

**3. Dietary sources of lutein:** Serving size is used for detecting the dietary sources of lutein. It is a standardized amount of food and used to quantify recommended amounts of food groups, or represent quantities that people typically consume on a Nutrition Facts label. One serving size is equal to: one medium sized egg, one cup of cooked spinach, one large green pepper and one cup of raw (tomato, leek, parsley and lettuce), one medium sized orange and one third cup or handful nuts (12). Recommended level for eye health: 10 milligrams (mg)/day for lutein and 2 mg/day for zeaxanthin (26). The lutein contents of one cup of Spinach, one egg yolk, half cup of parsley and one ounce of pistachios are containing about 20.4 mg (26), 0.1, 1.2 mg and 1.4 mg respectively (1, 27, 28).

**4. Anthropometric measures** (body weight, height) were measured by stadiometer, and the formula used to determine body mass index (BMI) was weight in kilograms divided by square height in meters. Class I obesity (BMI 30-34.9), class II obesity (BMI 35-39.9), class III obesity (BMI  $\geq 40$ ), underweight (BMI  $< 18.5$ ), normal weight (BMI = 18.5 -24.9), and overweight (BMI = 25-29.9) were the BMI classifications assigned to the patients (29-

32). Mid upper arm circumference was measured by tape measure (MUAC). Normal MUAC for adult are >23 cm for male and >22cm for women (33).

**5. Ophthalmic examination:** All patients were examined of visual acuity and intraocular pressure and senior ophthalmologist was examined posterior segment (vitreous and retina) using slit-lamp biomicroscopy.

**6. Optical coherence tomography (OCT) for macular assessment:** Is a non-invasive test that provides color-coded, cross sectional images of the retina to enable early detection and treatment of ocular disease that may develop without any noticeable symptoms. The OCT scan uses a low- coherence light to obtain higher resolution images of the layers of the retina and optic nerve. The color-coded images provide a wealth of information to measure the thickness of the retina and identify any optic nerve abnormalities (7).

**7. Measurement of the level of glycosylated hemoglobin (HbA1c):** was tested in laboratory of Imamein Kadhimein Medical City. The hemoglobin A1C (glycated hemoglobin, glycosylated hemoglobin, HbA1c) test is used to assess glucose control levels and diagnose diabetes. It is an average of the blood sugar level over for previous three months and represented in a percentage. Hemoglobin is a protein which only found in red blood cells. The main job of hemoglobin is to carry oxygen from the lungs to all the cells of the body. Hemoglobin becomes glycated or coated with glucose from the

bloodstream. The amount of glucose that is present in the blood will attach to the hemoglobin protein, and increased glucose levels will reflect on the surface of the hemoglobin protein, thereby making a higher HbA1c level (34). HbA1c test below 5.7 % classify as normal, or in the non-diabetic range, HbA1c value of 5.7 % to 6.4 % is considered to be pre-diabetic, while HbA1c of 6.5% or higher can be diagnosed as diabetes (35).

### Statistical Analysis

Data input and analysis were performed using the Statistical Package for Social Sciences, version 24 (SPSS 24) program. Simple frequency and percentage measures were used to display the data. The chi square test was used to determine the significance of the association between the variables, and Z test which is used for testing the significance of association between two proportions. A p value of 0.05 or less was deemed statistically significant.

### Results

#### 1- Age distribution:

This study showed 36% of diabetic patient had normal retina and 64% of them had diabetic retinopathy. Sex distribution was 41% male and 59% female. 50% of patients were with age 40-60 years and 49% were with age more than 60 years (Table 1).

**Table (1):** Frequency distribution of age of diabetic patients.

Age (year)	No.	%
< 40	1	1.0
40-60	50	50.0
> 60	49	49.0
Total	100	100.0

**2- Body mass index of diabetic patients:**

Thirtyeight percent of patients were with class I obesity, 12% with class II and 7% with class

III. While 24% of patients were with normal weight and 19% with overweight (Table 2).

**Table (2):** Frequency distribution of body mass index (BMI) of diabetic patients.

BMI	No.	%
Normal weight	24	24.0
Over weight	19	19.0
Obesity class I	38	38.0
Obesity class II	12	12.0
Obesity class III	7	7.0
Total	100	100.0

**3- Diabetic retinal lesions and lutein rich diet consumption:**

It is about 49% of diabetic patients had diabetic macular edema, 10% with non-proliferative diabetic retinopathy, 5%

with proliferative diabetic retinopathy and 36% of patients had normal retina (Table 3).

**Table (3):** Frequency distribution of diabetic retinal lesions in diabetic patients.

Retinal lesion	No.	%
Normal retina	36	36.0
Non proliferative diabetic retinopathy	10	10.0
Diabetic macular edema	49	49.0
Proliferative diabetic retinopathy	5	5.0
Total	100	100.0

The proportion of diabetic patients with normal retina which consume tomato, egg, green pepper, basil, spinach, parsley, leek, lettuce, orange and nut were 91.7%, 75%, 63.9%, 63.9%, 61.1%, 61.1%, 58.3, 52.8%, 30.6%, 33.3% respectively from total number of patients with normal retina, while the proportion of diabetic patients with retinopathy which consume tomato, egg, green pepper, basil, spinach, parsley, leek, lettuce, orange and nut were 76.6%, 46.9%, 32.8%, 9.4%, 28.1%, 20.3%, 7.8%, 4.7%,

14%, 10.9 respectively with significant p-values (0.004, 0.005, 0.003, 0.0001, 0.001, 0.0001, 0.0001, 0.0001, 0.04, 0.007) respectively. The proportion of diabetic patients with normal retina that consume 1-2 serving/week and 3-7 serving/week pistachio are 30.6% and 25% respectively, while the proportion of diabetic patients with retinopathy which consume 1-2 serving/week and 3-7 serving/week pistachio are 3.1% and 9.4% with significant p-value (0.0001, 0.03) respectively (Table 4).

**Table (4):** Proportion of lutein rich diet consumption among diabetic patients.

Type of Food	No. of serving/week	Normal retina, n=36		Diabetic retinopathy, n=64		P value	Total
		No.	Proportion	No.	Proportion		
Egg	7-35	27	75	30	46.9	0.005*	100
Spinach	1-2	22	61.1	18	28.1	0.001*	
Green pepper	3-7	23	63.9	21	32.8	0.003*	
Tomato	3-7	33	91.7	49	76.6	0.004*	
Leek	3-7	21	58.3	5	7.8	0.0001*	
Parsley	3-7	22	61.1	13	20.3	0.0001*	
Lettuce	3-7	19	52.8	3	4.7	0.0001*	
Basil	3-7	23	63.9	6	9.4	0.0001*	
Orange	3-7	11	30.6	9	14	0.04*	
Nut	3-7	12	33.3	7	10.9	0.007*	
Pistachio	1-2	11	30.6	2	3.1	0.0001*	
	3-7	9	25	6	9.4	0.03*	

It is about 91.7% of diabetic patients with normal retina were consume 3-7 serving/week tomato, 75% were consume 7-35 serving/week egg, 75% were consume 1-7 serving/week zucchini, 72% were consume 1-3 serving/week spinach, 63.9% were consume 3-7 serving/week green pepper, 63.9% were consume 3-7 serving/week basil, 61% were consume 3-7 serving/week parsley, 58% were consume 3-7 serving/week leek, 52.8% were consume 3-7 serving/week lettuce, 41.7%

were consume 1-2 serving/week nut and 30.6% were consume 1-2 serving/week pistachio, while 76.6% of diabetic patients with retinopathy were consume 3-7 serving/week tomato, 40.9% were consume 7-35 serving/week egg, 51.6% were consume 1-7 serving/week zucchini, 31% were consume 1-3 serving/week spinach, 40.6% were consume 1-2 serving/week green pepper, 14% were consume 1-2 serving/week basil, 29.7% were consume 1-2 serving/week parsley,

12.5% were consume 1-2 serving/week leek, 17% were consume 1-2 serving/week lettuce, 21.9% were consume 1-2 serving/week nut and 3% were consume 1-2 serving/week

pistachio with significant p-value (0.01, 0.006, 0.02, 0,0001, 0.005, 0.0001, 0.0001, 0,0001, 0.0001, 0.0001, 0.0001) respectively (Table 5).

**Table 5:** Frequency distribution of lutein rich diet consumption among diabetic patients.

Type of Food	Normal retina, n=36				Diabetic retinopathy, n=64				P value	Total
	No serving	Serving			No serving	Serving				
		Serving /week	No.	%		Serving /week	No.	%		
Egg	4	1-6	5	13.9	8	1-6	26	40.6	0.006*	100
		7-35	27	75		7-35	30	40.9		
Spinach	10	1-3	26	72	44	1-3	20	31	0.0001*	
Zucchini	9	1-7	27	75	31	1-7	33	51.6	0.02*	
Green pepper	5	1-2	8	22	17	1-2	26	40.6	0.005*	
		3-7	23	63.9		3-7	21	32.8		
Tomato	1	1-2	2	5.6	5	1-2	10	15.6	0.01*	
		3-7	33	91.7		3-7	49	76.6		
Leek	6	1-2	9	25	51	1-2	8	12.5	0.0001*	
		3-7	21	58		3-7	5	7.8		
Parsley	6	1-2	8	22	32	1-2	19	29.7	0.0001*	
		3-7	22	61		3-7	13	20		
Lettuce	5	1-2	12	33	50	1-2	11	17	0.0001*	
		3-7	19	52.8		3-7	3	4.7		
Basil	6	1-2	7	19	49	1-2	9	14	0.0001*	
		3-7	23	63.9		3-7	6	9.4		
Nut	9	1-2	15	41.7	43	1-2	14	21.9	0.0001*	
		3-7	12	33		3-7	7	10.9		
Pistachio	16	1-2	11	30.6	56	1-2	2	3	0.0001*	
		3-7	9	25		3-7	6	9.3		

## Discussion

Diabetes mellitus is a group of metabolic diseases that all have a hyperglycemic phenotype, and may associated with obesity which is a major public health problem worldwide (36- 38). Visual impairment is a global issue, particularly in developing nations and among the most common causes of avoidable blindness and moderate-to-severe visual impairment from 1990 to 2020, diabetic retinopathy ranks fifth globally. It is also one of the main causes of vision loss (9, 39). Common carotenoid pigments include lutein and zeaxanthin, which are present in

high concentrations in egg yolks, orange and yellow fruits, and dark green vegetables. Because of its antioxidant qualities, position inside the retina, and capacity to absorb oxidative blue light, lutein may offer protection against DR (2). Age-related macular degeneration (AMD) and DR are treated and prevented with dietary and lifestyle changes (40). The current study showed that 75% of diabetes individuals with normal retinas consume 7–35 servings of eggs per week, and this intake was substantially correlated with normal retinal exams in these patients. Previous studies showed that egg

lutein content has been linked to eye health, the egg yolk provides an excellent dietary source of lutein and zeaxanthin because the bioavailability from the yolk matrix is much higher than from the leaves of green vegetables. The high bioavailability of a fat-soluble nutrient such as zeaxanthin from the egg is due to the rich lipid matrix of the yolk. Egg yolk is a good dietary source of both zeaxanthin and lutein, particularly as part of a typical western diet, which is poor in vegetables and fruits. A high intake of lutein can also increase the macular content of meso-zeaxanthin because the lutein can convert to meso-zeaxanthin in the central retina (1, 26, 27). In the current study consuming spinach, zucchini, green pepper, tomato, leek, parsley, lettuce, basil, oranges, nuts, and pistachios is significantly linked to normal retina function. A weekly consumption of 1-2 serving\week (1-2 cups) of cooked spinach was reported by 61.1% of patients with normal retinal examinations. Previous studies showed that lutein and zeaxanthin are the most common xanthophylls in green leafy vegetables like spinach which had important role in eye health (1). In this study, 63.9% of patients consumed 3-7 servings (one and a half cups) of chopped green pepper or one large bell pepper were with normal retina. Peppers are one of the most widely consumed foods throughout the world owing to their attractive colors and strong flavor, green peppers get their unique yellow and orange hue from lutein, a chemical substance that's abundant in them. Research has demonstrated that lutein, an antioxidant, enhances eye health (41, 42). Carotenoids act as antioxidants and deactivating free radicals. Nutritional supply of carotenoids with ocular health benefits such as lutein and zeaxanthin,

with potential health benefits for humans (41, 42). Each week, about 91.7% of people consumed three to seven servings (one cup) of chopped or sliced fresh tomatoes. Lutein is becoming increasingly important in preventive medicine due to its possible role in maintaining good vision. Research has demonstrated that a unique pigment found in tomatoes called lutein serves as a filter to shield the eyes from harmful light and oxygen which showed higher antioxidant capacity that protect the retina and lens against ultraviolet and blue spectrum light. Study in Panama City at 2017 showed that the most consumed lutein and zeaxanthin food sources in the study population were tomatoes, egg yolks and green peppers. Lutein and zeaxanthin, both oxygen-containing carotenoids in tomatoes and tomato-based food products are considered to play vital roles in promoting ocular development and maintaining eye health (28, 43- 45). Among the study group of diabetes patients, those who consumed 3-7 servings (one cup) of raw leek per week accounted for 58% of normal retinal tissue. Previous studies showed significant eye problems can be avoided and maintained with the use of leafy green vegetables like leek. Lutein and zeaxanthin are dietary carotenoids derived from dark green leafy vegetables, orange and yellow fruits that form the macular pigment of the human eyes. It was assumed that they protect against visual disorders such as hypoxia induced retinopathy and diabetic retinopathy. The mechanism by which they are involved in the prevention of eye diseases may be due their local antioxidant activity and physical blue light filtration properties which are the direct biological effects of lutein and also improve normal ocular function by



enhancing contrast sensitivity and by reducing glare disability which is the photophobia and discomfort when intense light enters the eye (45- 50). Approximately 61.1% of diabetes patients with normal retinal function consumed three to seven servings (one cup of three-quarters bunch or one tablespoon of freshly chopped parsley) of parsley each week. Several studies documented that consumption of dark green leafy vegetables had a protective role against eye diseases. Parsley is an important culinary herb originated from the Mediterranean region. Its main constituent's carotenoids, flavonoids and it has anti oxidative activity (1, 51- 53).

Five to seven servings (two cups of raw chopped lettuce) were consumed weekly by 52.8% of the diabetic individuals in this study with normal retinal results. Numerous studies have identified lutein and zeaxanthin to be essential components for eye health, they constitute the main pigments found in the yellow spot of the human retina which protect the macula from damage by blue light, improve visual acuity and remove harmful reactive oxygen species. Parsley and lettuce are rich in antioxidants, lutein and zeaxanthin, which help shield the surface of the eyes and prevent eye diseases. A study in the US showed that a higher dietary intake of carotenoids, specifically lutein is associated with reduced eye diseases (1, 51- 53). In the current study, a weekly consumption of 3–7 servings (one cup fresh or 2 tablespoons chopped) of basil was reported by about 63.9% of normal retinal examinations. Basil supports eye health, and fights free radicals through its antioxidant and anti-inflammatory properties which is high in beta carotene, lutein and zeaxanthin (54- 57). About 30.6%

of diabetic individuals with normal retinal examinations consumed oranges, consuming three to seven servings (one medium-sized piece) per week. Nutrition plays a vital role in human health with no exception to the eye. Healthy eyes provide good vision, which is essential for an enjoyable and productive lifestyle. Numerous studies have identified lutein and zeaxanthin to be essential components for eye health. Consumption of orange have protective effect on the retina. Orange is rich with lutein and zeaxanthin which have a strong antioxidant content (1, 18). Approximately 33.3% of diabetes patients with normal retinal examinations consumed 3–7 servings (equivalent to 1/3 of a cup or one handful) of nuts each week. In diabetes individuals, around 30.6% of normal retina were consumed in 1-2 servings (one handful or ten pieces) per week, while 25% consumed 3-7 servings per week. Of all nuts, pistachios contain the highest concentrations of zeaxanthin and lutein. These nutrients guard your eyes against macular degeneration, an eye condition that can cause vision loss as you age, and blue light damage (19, 21).

Other foods high in lutein, such as kale, which is not available in Iraq, collard greens, squash, broccoli, peas, orange pepper, corn, persimmon, tangerines, carrots, kiwis, grapes, and bananas, were not significantly associated with a protective effect on the retina. This could be because our patients did not have access to these foods or because their prices made them difficult to purchase in addition to their poor dentation.

## Conclusions

Patients with diabetes can avoid retinal damage by consuming the recommended weekly servings of lutein-rich foods

especially tomato, eggs, zucchini, spinach, green pepper, basil, parsley, leek, lettuce, nut and pistachio.

### Recommendations

1. Suggested daily consumption of lutein rich foods for diabetic patients.
2. Encourage weight reduction.
3. Increase knowledge about types of lutein rich foods.
4. Routine ophthalmic examination.

**Source of Funding:** The current study was funded by our charges with no any other funding sources elsewhere.

**Ethical Clearance:** Imamein Kadhimein Medical City and the Arabic Council of Medical Specialization had granted their official approval. After informing the patients about the purpose and goals of the study, assuring their privacy, and ensuring that the questionnaires were completed anonymously, the patients gave their informed consent. (Document no. 2024HRS872).

**Conflict of Interest:** Non

### References

1. Abdel-Aal el-SM, Akhtar H, Zaheer K, Ali R. Dietary sources of lutein and zeaxanthin carotenoids and their role in eye health. *Nutrients*. 2013 Apr 9; 5(4):1169-1185.

Doi: [10.3390/nu5041169](https://doi.org/10.3390/nu5041169). [PubMed]

2. Sahli MW, Mares JA, Meyers KJ, Klein R, Brady WE, Klein BE, Ochs-Balcom HM, Donahue RP, Millen AE. Dietary Intake of Lutein and Diabetic Retinopathy in the Atherosclerosis Risk in Communities Study (ARIC). *Ophthalmic Epidemiology*. 2016; 23(2):99-108.

Doi: [10.3109/09286586.2015.1129426](https://doi.org/10.3109/09286586.2015.1129426). [PubMed]

3. Abusaib, Mohammed et al. Iraqi Experts Consensus on the Management of Type 2 Diabetes/ Prediabetes in Adults. *Clinical Medicine Insights Endocrinology and Diabetes*. 2020 Aug 19; 13: 1179551420942232.

Doi: [10.1177/1179551420942232](https://doi.org/10.1177/1179551420942232). [PubMed]

4. Al Ashoor M, Al Hamza A, Zaboon I, Almomin A, Mansour A. Prevalence and risk factors of diabetic retinopathy in Basrah, Iraq. *J Med Life*. 2023 Feb; 16(2):299-306.

Doi: [10.25122/jml-2022-0170](https://doi.org/10.25122/jml-2022-0170). [PubMed]

5. Teo ZL, Tham YC, Yu M, et al. Global Prevalence of Diabetic Retinopathy and Projection of Burden through 2045: Systematic Review and Meta-analysis. *Ophthalmology*. 2021 Nov; 128(11):1580-1591.

Doi: [10.1016/j.ophtha.2021.04.027](https://doi.org/10.1016/j.ophtha.2021.04.027). [PubMed]

6. Shah J, Cheong ZY, Tan B, Wong D, Liu X, Chua J. Dietary Intake and Diabetic Retinopathy: A Systematic Review of the Literature. *Nutrients*. 2022; 14(23):5021.

Doi: [10.3390/nu14235021](https://doi.org/10.3390/nu14235021). [PMC]

7. Basic and Clinical Science Course. 2021. Retina and Vitreous. In: American Academy of Ophthalmology. *Fundamentals and Principles of Ophthalmology*. San Francisco, CA: American Academy of Ophthalmology; p. (91-93) and (25-28).

8. Bowling Brad. 2016. Kanski's Clinical Ophthalmology (A systematic approach); Eighth edition. P: 521 and 522.

9. GBD 2019 Blindness and Vision Impairment Collaborators; Vision Loss

Expert Group of the Global Burden of Disease Study. Causes of blindness and vision impairment in 2020 and trends over 30 years, and prevalence of avoidable blindness in relation to VISION 2020: the Right to Sight: an analysis for the Global Burden of Disease Study. *Lancet Glob Health*. 2021 Feb; 9(2):e144-e160.

Doi: [10.1016/S2214-109X\(20\)30489-7](https://doi.org/10.1016/S2214-109X(20)30489-7).  
[Google Scholar]

10.Dow C, Mancini F, Rajaobelina K, Boutron-Ruault MC, Balkau B, Bonnet F, Fagherazzi G. Diet and risk of diabetic retinopathy: a systematic review. *European journal of epidemiology*. 2018 Feb; 33(2):141-156.

Doi: [10.1007/s10654-017-0338-8](https://doi.org/10.1007/s10654-017-0338-8).  
[PubMed]

11.Yau JW, Rogers SL, Kawasaki R, et al. Global prevalence and major risk factors of diabetic retinopathy. *Diabetes Care*. 2012; 35(3):556-564.

Doi: [10.2337/dc11-1909](https://doi.org/10.2337/dc11-1909). [PubMed]

12.Raymond Janice, Morrow Kelly. 2023. Krause and Mahan's Food and the Nutrition Care Process; 16th Edition. P (655-656) and (1114 – 1126).

13.Moschos MM, Dettoraki M, Tsatsos M, Kitsos G, Kalogeropoulos C. Effect of carotenoids dietary supplementation on macular function in diabetic patients. *Eye and vision (London)*. 2017 Oct 15; 4: 23.

Doi: [10.1186/s40662-017-0088-4](https://doi.org/10.1186/s40662-017-0088-4).  
[PubMed]

14.Buscemi S, Corleo D, Di Pace F, Petroni ML, Satriano A, Marchesini G. The Effect of Lutein on Eye and Extra-Eye Health. *Nutrients*. 2018 Sep 18; 10(9):1321.

Doi: [10.3390/nu10091321](https://doi.org/10.3390/nu10091321). [PubMed]

15.Ahn YJ, Kim H. Lutein as a Modulator of Oxidative Stress-Mediated Inflammatory Diseases. *Antioxidants (Basel)*. 2021 Sep 13; 10(9):1448.

Doi:[10.3390/antiox10091448](https://doi.org/10.3390/antiox10091448).  
[PubMed]

16.Li LH, Lee JC, Leung HH, Lam WC, Fu Z, Lo ACY. Lutein Supplementation for Eye Diseases. *Nutrients*. 2020 Jun 9; 12(6):1721.

Doi: [10.3390/nu12061721](https://doi.org/10.3390/nu12061721). [PubMed]

17.Neelam K, Goenadi CJ, Lun K, Yip CC, Au Eong KG. Putative protective role of lutein and zeaxanthin in diabetic retinopathy. *The British journal of ophthalmology*. 2017 May; 101(5):551-558.

Doi: [10.1136/bjophthalmol-2016-309814](https://doi.org/10.1136/bjophthalmol-2016-309814). [PubMed]

18.Bernstein PS, Li B, Vachali PP, Gorusupudi A, Shyam R, Henriksen BS, Nolan JM. Lutein, zeaxanthin, and meso-zeaxanthin: The basic and clinical science underlying carotenoid-based nutritional interventions against ocular disease. *Progress in retinal and eye research*. 2016 Jan; 50:34-66.

Doi: [10.1016/j.preteyeres.2015.10.003](https://doi.org/10.1016/j.preteyeres.2015.10.003).  
[PubMed]

19.Li X, Holt RR, Keen CL, Morse LS, Zivkovic AM, Yiu G, Hackman RM. Potential roles of dietary zeaxanthin and lutein in macular health and function. *Nutrition reviews*. 2023 May 10; 81(6):670-683.

Doi: [10.1093/nutrit/nuac076](https://doi.org/10.1093/nutrit/nuac076). [PubMed]

20.Eisenhauer B, Natoli S, Liew G, Flood VM. Lutein and Zeaxanthin-Food Sources, Bioavailability

- and Dietary Variety in Age-Related Macular Degeneration Protection. *Nutrients*. 2017 Feb 9; 9(2):120.  
[Doi: 10.3390/nu9020120](https://doi.org/10.3390/nu9020120). [[PubMed](#)]
- 21.Mrowicka M, Mrowicki J, Kucharska E, Majsterek I. Lutein and Zeaxanthin and Their Roles in Age-Related Macular Degeneration-Neurodegenerative Disease. *Nutrients*. 2022 Feb 16; 14(4):827.  
[Doi: 10.3390/nu14040827](https://doi.org/10.3390/nu14040827). [[PubMed](#)]
- 22.Wong MYZ, Man REK, Fenwick EK, Gupta P, Li LJ, van Dam RM, Chong MF, Lamoureux EL. Dietary intake and diabetic retinopathy: A systematic review. *PLoS One*. 2018 Jan 11; 13(1):e0186582.  
[Doi: 10.1371/journal.pone.0186582](https://doi.org/10.1371/journal.pone.0186582). [[Google Scholar](#)]
- 23.Sala-Vila A, Díaz-López A, Valls-Pedret C, et al. Dietary Marine  $\omega$ -3 Fatty Acids and Incident Sight-Threatening Retinopathy in Middle-Aged and Older Individuals With Type 2 Diabetes: Prospective Investigation From the PREDIMED Trial. *JAMA Ophthalmology*. 2016 Oct 1; 134(10):1142-1149.  
[Doi:10.1001/jamaophthalmol.2016.2906](https://doi.org/10.1001/jamaophthalmol.2016.2906). [[Google Scholar](#)]
- 24.Chua J, Chia A-R, Chee ML, Man REK, Tan GSW, Lamoureux EL, Wong TY, Chong MF-F, Schmetterer L.. The Relationship of Dietary Fish Intake and Diabetic Retinopathy in Asian Patients with Type 2 Diabetes. *Investigative Ophthalmology & Visual Science*. 2017 June; 58(8) 4284.  
[Doi: 10.1038/s41598-017-18930-6](https://doi.org/10.1038/s41598-017-18930-6). [[Google Scholar](#)]
- 25.Omer W, Al-Hadithi T. Developing a socioeconomic index for health research in Iraq. *Eastern Mediterranean health journal*. 2017 Dec 14; 23(10):670-677.  
[DOI:10.26719/2017.23.10.670](https://doi.org/10.26719/2017.23.10.670). [[PubMed](#)]
- 26.Widomska J, SanGiovanni JP, Subczynski WK. Why is Zeaxanthin the Most Concentrated Xanthophyll in the Central Fovea? *Nutrients*. 2020 May 7; 12(5):1333.  
[Doi: 10.3390/nu12051333](https://doi.org/10.3390/nu12051333). [[PubMed](#)]
- 27.Mares J. Lutein and Zeaxanthin Isomers in Eye Health and Disease. *Annual review of nutrition*. 2016 Jul 17; 36:571-602.  
[Doi: 10.1146/annurev-nutr-071715-051110](https://doi.org/10.1146/annurev-nutr-071715-051110). [[PubMed](#)]
- 28.Giorio G, Yildirim A, Stigliani AL, D'Ambrosio C. Elevation of lutein content in tomato: a biochemical tug-of-war between lycopene cyclases. *Metabolic engineering*. 2013 Nov; 20:167-176.  
[Doi: 10.1016/j.ymben.2013.10.007](https://doi.org/10.1016/j.ymben.2013.10.007). [[PubMed](#)]
- 29.World Health Organization. *Obesity: Preventing and Managing the Global Epidemic*. Volume 894 WHO; Geneva, Switzerland: 2000. WHO Consultation on Obesity (1999: Geneva, Switzerland) Report of a WHO Consultation. World Health Organization Technical Report Series. [[Google Scholar](#)]
- 30.Kadhom EH and Radhi NJ. The relation between oral health and body mass index among women with hyperthyroidism. *Diyala Journal of Medicine* 30 October 2023; 25 (1): 79-87  
[Doi: 10.26505/DJM.25017220105](https://doi.org/10.26505/DJM.25017220105).

31. Abdullah AA and Zangana S N. Correlation between body mass index and in-hospital mortality in patients with ST-segment elevation myocardial infarction in Erbil city- Iraq. *Diyala Journal of Medicine* 25 October 2021; 21(1):35-43  
Doi: [10.26505/DJM.21015203922](https://doi.org/10.26505/DJM.21015203922).
32. Mubarak AA and Mustaf AH. Influence of Body Mass Index (BMI) on Outcome of Colon Cancer in Relationships with Other Clinicopathological Factors. *Diyala Journal of Medicine* 25 December 2021; 21(2): 52-63.  
Doi: [10.26505/DJM.21026060526](https://doi.org/10.26505/DJM.21026060526).
33. Yallamraju SR, Mehrotra R, Sinha A, Gattumedi SR, Gupta A, Khadse SV. Use of mid upper arm circumference for evaluation of nutritional status of OSMF patients. *Journal of International Society of Preventive & Community Dentistry*. 2014 Dec; 4(Suppl 2):S122-125.  
Doi: [10.4103/2231-0762.146217](https://doi.org/10.4103/2231-0762.146217).  
[PubMed]
34. Al-Ansary L, Farmer A, Hirst J, Roberts N, Glasziou P, Perera R, Price CP. Point-of-care testing for Hb A1c in the management of diabetes: a systematic review and metaanalysis. *Clinical Chemistry*. 2011 Apr; 57(4): 568-576.  
Doi: [10.1373/clinchem.2010.157586](https://doi.org/10.1373/clinchem.2010.157586).  
[PubMed]
35. Sikaris KA. The role and quality of Hb A1C: a continuing evolution. *Clinical Chemistry*. 2015 May; 61(5): 689-690.  
Doi: [10.1373/clinchem.2015.239319](https://doi.org/10.1373/clinchem.2015.239319).  
[PubMed]
36. Raqib TM, Polus RK and Mohammad NS. Prevalence of Vitamin B12 Deficiency in Patients with type 2 Diabetes Mellitus on Metformin. *Diyala Journal of Medicine* 15 October 2022; 23(1): 22-32.  
Doi: [10.26505/DJM.23016480407](https://doi.org/10.26505/DJM.23016480407).
37. Salih KKH, Ali SJ and Ahmed MAA. Prevalence of Obesity among Students in Private and Public high Schools in Sulaimani City. *Diyala Journal of Medicine* 30 June 2023; 24(2): 24-3  
Doi: [10.26505/DJM.24026991016](https://doi.org/10.26505/DJM.24026991016).
38. Ibrahim GI, et al. Association between Serum Levels of Vitamin D, Vitamin B12 and Folate with Oxidative Biomarkers in Diabetic Type 2 Patients in People in Erbil City: A Case-Control Study. *Diyala Journal of Medicine* december 2020; 19(2): 200-207.  
Doi: [10.26505/DJM.19025660915](https://doi.org/10.26505/DJM.19025660915).
39. Saeed HR, Dadoosh AG, Ali BM and Elbassiouny KAM. The Association between Vitamin D3 Deficiency and Cataract Formation in Baghdad Al-Karkh. *Diyala Journal of Medicine* 25 October 2024; 27 (1): 35-49  
Doi: [10.26505/DJM.27018630624](https://doi.org/10.26505/DJM.27018630624)
40. Rondanelli M, Gasparri C, Riva A, Petrangolini G, Barrile GC, Cavioni A, Razza C, Tartara A, Perna S. Diet and ideal food pyramid to prevent or support the treatment of diabetic retinopathy, age-related macular degeneration, and cataracts. *Frontiers in medicine (Lausanne)*. 2023 May 30; 10:1168560.  
Doi: [10.3389/fmed.2023.1168560](https://doi.org/10.3389/fmed.2023.1168560).  
[PubMed]
41. Rodríguez-Rodríguez E, Sánchez-Prieto M, Olmedilla-Alonso B. Assessment of carotenoid concentrations in red peppers (*Capsicum annuum*) under domestic refrigeration for three weeks as determined by HPLC-DAD. *Food chemistry: X*. 2020 May 28; 6:100092.  
Doi: [10.1016/j.fochx.2020.100092](https://doi.org/10.1016/j.fochx.2020.100092).  
[PubMed]

42. Xu J, Lin J, Peng S, Zhao H, Wang Y, Rao L, Liao X, Zhao L. Development of an HPLC-PDA Method for the Determination of Capsanthin, Zeaxanthin, Lutein,  $\beta$ -Cryptoxanthin and  $\beta$ -Carotene Simultaneously in Chili Peppers and Products. *Molecules*. 2023 Mar 3; 28(5):2362.  
Doi: [10.3390/molecules28052362](https://doi.org/10.3390/molecules28052362). [PubMed]
43. Wu Y, Yuan Y, Jiang W, Zhang X, Ren S, Wang H, Zhang X, Zhang Y. Enrichment of health-promoting lutein and zeaxanthin in tomato fruit through metabolic engineering. *Synth Syst Biotechnol*. 2022 Aug 24; 7(4):1159-1166.  
Doi: [10.1016/j.synbio.2022.08.005](https://doi.org/10.1016/j.synbio.2022.08.005). [PubMed]
44. Alvarado-Ramos KE, De Leon L, Fontes F, Rios-Castillo I. Dietary Consumption of Lutein and Zeaxanthin in Panama: A Cross-Sectional Study. *Current developments in nutrition*. 2018 Aug 6; 2(9): nzy064.  
Doi: [10.1093/cdn/nzy064](https://doi.org/10.1093/cdn/nzy064). [PubMed]
45. Khachik F, Carvalho L, Bernstein PS, Muir GJ, Zhao DY, Katz NB. Chemistry, distribution, and metabolism of tomato carotenoids and their impact on human health. *Experimental biology and medicine* (Maywood). 2002 Nov; 227(10):845-851.  
Doi: [10.1177/153537020222701002](https://doi.org/10.1177/153537020222701002). [PubMed]
46. Jia YP, Sun L, Yu HS, Liang LP, Li W, Ding H, Song XB, Zhang LJ. The Pharmacological Effects of Lutein and Zeaxanthin on Visual Disorders and Cognition Diseases. *Molecules*. 2017 Apr 20; 22(4):610.  
Doi: [10.3390/molecules22040610](https://doi.org/10.3390/molecules22040610). [PubMed]
47. Ma L, Lin XM. Effects of lutein and zeaxanthin on aspects of eye health. *Journal of the science of food and agriculture*. 2010 Jan 15; 90(1):2-12.  
Doi: [10.1002/jsfa.3785](https://doi.org/10.1002/jsfa.3785). PMID: 20355006. [PubMed]
48. Sommerburg O, Keunen JE, Bird AC, van Kuijk FJ. Fruits and vegetables that are sources for lutein and zeaxanthin: the macular pigment in human eyes. *The British journal of ophthalmology*. 1998 Aug; 82(8):907-10.  
Doi: [10.1136/bjo.82.8.907](https://doi.org/10.1136/bjo.82.8.907). [PubMed]
49. Ahn YJ, Kim H. Lutein as a Modulator of Oxidative Stress-Mediated Inflammatory Diseases. *Antioxidants* (Basel). 2021 Sep 13; 10(9):1448.  
Doi: [10.3390/antiox10091448](https://doi.org/10.3390/antiox10091448). [PubMed]
50. Walsh RP, Bartlett H, Eperjesi F. Variation in Carotenoid Content of Kale and Other Vegetables: A Review of Pre- and Post-harvest Effects. *Journal of agricultural and food chemistry*. 2015 Nov 11; 63(44):9677-9682.  
Doi: [10.1021/acs.jafc.5b03691](https://doi.org/10.1021/acs.jafc.5b03691). [PubMed]
51. Mahmood S, Hussain S, Malik F. Critique of medicinal conspicuousness of Parsley (*Petroselinum crispum*): a culinary herb of Mediterranean region. *Pakistan journal of pharmaceutical sciences*. 2014 Jan; 27(1):193-202. PMID: 24374449. [PubMed]
52. de Oliveira VS, Chávez DWH, Paiva PRF, Gamallo OD, Castro RN, Sawaya

ACHF, Sampaio GR, Torres EAFDS, Saldanha T. Parsley (*Petroselinum crispum* Mill.): A source of bioactive compounds as a domestic strategy to minimize cholesterol oxidation during the thermal preparation of omelets. *Food research international*. 2022 Jun; 156:111199.

Doi: [10.1016/j.foodres.2022.111199](https://doi.org/10.1016/j.foodres.2022.111199).  
[PubMed]

53.Mrowicka M, Mrowicki J, Kucharska E, Majsterek I. Lutein and Zeaxanthin and Their Roles in Age-Related Macular Degeneration-Neurodegenerative Disease. *Nutrients*. 2022 Feb 16; 14(4):827.

Doi: [10.3390/nu14040827](https://doi.org/10.3390/nu14040827). [PubMed]

54.Romano R, De Luca L, Aiello A, Pagano R, Di Pierro P, Pizzolongo F, Masi P. Basil (*Ocimum basilicum* L.) Leaves as a Source of Bioactive Compounds. *Foods*. 2022 Oct 14; 11(20):3212.

Doi: [10.3390/foods11203212](https://doi.org/10.3390/foods11203212). [PubMed]

55.Vassilina G, Sabitova A, Idrisheva Z, Zhumabekova A, Kanapiyeva F, Orynassar R, Zhamanbayeva M, Kamalova M, Assilbayeva J, Turgumbayeva A, Abilkassymova A. Bio-active compounds and major biomedical properties of basil (*Ocimum basilicum*, lamiaceae). *Nat Prod Res*. 2024 May 30:1-19.

Doi: [10.1080/14786419.2024.2357662](https://doi.org/10.1080/14786419.2024.2357662).  
[PubMed]

56.Sikora M, Złotek U, Kordowska-Wiater M, Świeca M. Effect of Basil

Leaves and Wheat Bran Water Extracts on Antioxidant Capacity, Sensory Properties and Microbiological Quality of Shredded Iceberg Lettuce during Storage. *Antioxidants (Basel)*. 2020 Apr 24; 9(4):355.

Doi: [10.3390/antiox9040355](https://doi.org/10.3390/antiox9040355). [PubMed]

57.Hakkim FL, Shankar CG, Girija S. Chemical composition and antioxidant property of holy basil (*Ocimum sanctum* L.) leaves, stems, and inflorescence and their in vitro callus cultures. *Journal of agricultural and food chemistry*. 2007 Oct 31; 55(22):9109-9117.

Doi: [10.1021/jf071509h](https://doi.org/10.1021/jf071509h). [PubMed]

## تأثير تناول الأطعمة الغنية باللوتين على الشبكية وارتباطه باعتلال الشبكية لدى مرضى السكري

هبة رعد سعيد<sup>٢</sup>, احمد غازي دعدوش<sup>٣</sup>, بسمه محمد علي<sup>٤</sup>, خالد عوض محمد البسيوني

### الملخص

**خلفية الدراسة:** اعتلال الشبكية السكري هو نتيجة شائعة للأوعية الدموية الدقيقة لمرض السكري الذي يؤثر على العينين. إذا لم يتم علاجه، يتسبب اعتلال الشبكية السكري في تلف الأوعية الدموية في شبكية العين ويؤدي إلى العمى أو فقدان البصر. يؤدي ارتفاع السكر في الدم على مدى فترة طويلة من الزمن إلى زيادة الإجهاد التأكسدي والتهابي ومسارات بروتين كيناز C التي تؤدي في النهاية إلى تلف بطانة الأوعية الدموية في شبكية العين وفقدان الحويصلة. اللوتين عبارة عن كاروتين ذو خصائص مضادة للالتهابات ومضادة للأكسدة.

**اهداف الدراسة:** لتحديد الأدوار الوقائية للتناول الغذائي للأطعمة الغنية باللوتين على شبكية العين وارتباطها باعتلال الشبكية لدى مرضى السكري.

**الحالات و المنهجية:** أجريت دراسة مقطعية في قسم طب العيون في مدينة الإمامين الكاظمين الطبية، بغداد، العراق، في الفترة من يوليو إلى نوفمبر ٢٠٢٣، وشملت ما مجموعه ١٠٠ مريض بالسكري. تم استخدام قائمة الاستبيانات التي تتكون من المعلومات الاجتماعية والديموغرافية والحالة الاجتماعية والاقتصادية والموارد الغذائية للوتين والقياسات البشرية وفحص العيون والتحقيقات التي تشمل التصوير المقطعي التوافقي البصري لتقييم البقعة الصفراء وتقييم نسبة HbA1c.

**النتائج:** أظهرت الدراسة الحالية أن ٦٤٪ من مرضى السكري يعانون من اعتلال الشبكية و ٣٦٪ لديهم شبكية طبيعية. أظهرت الدراسة أن ٩١,٧٪ من مرضى السكري ذوي الشبكية الطبيعية كانوا يتناولون ٧-٣ حصص/أسبوع من الطماطم، و ٧٥٪ كانوا يستهلكون ٣٥-٧ حصة/أسبوع من البيض، و ٧٥٪ كانوا يستهلكون ٧-١ حصص/أسبوع من الكوسا، و ٧٢٪ كانوا يستهلكون ١-٣ حصص/أسبوع من السبانخ، و ٦٣,٩٪ يستهلكون ٧-٣ حصص/أسبوع من الفلفل الأخضر، و ٦٣,٩٪ يستهلكون ٧-٣ حصص/أسبوع من الريحان، و ٦١٪ كانوا يستهلكون ٧-٣ حصص/أسبوع من البقدونس، و ٥٨٪ يستهلكون ٧-٣ حصص/أسبوع من الكراث، و ٥٢,٨٪ يستهلكون ٧-٣ حصص/أسبوع من الخس، و ٤١,٧٪ يستهلكون ٢-١ حصة/أسبوع من المكسرات و ٣٠,٦٪ يستهلكون ٢-١ حصة/أسبوع من الفستق مع القيمة الاحتمالية المعتمد بها احصائيا.

**الاستنتاجات:** يمكن لمرضى السكري تجنب تلف الشبكية عن طريق تناول الحصص الأسبوعية الموصى بها من الأطعمة الغنية باللوتين وخاصة الطماطم والبيض والكوسا والسبانخ والفلفل الأخضر والريحان والبقدونس والكراث والخس والمكسرات والفستق.

**الكلمات المفتاحية:** اعتلال الشبكية السكري، اللوتين، ضعف البصر.

البريد الإلكتروني: [hebars81@yahoo.com](mailto:hebars81@yahoo.com)

تاريخ استلام البحث: ١٧ تموز ٢٠٢٤

تاريخ قبول البحث: ٣ كانون الاول ٢٠٢٤

<sup>١</sup> شعبة السيطرة على الامراض الانتقالية/ قسم الصحة العامة/ دائرة صحة بغداد/ الرصافة/ العراق.

<sup>٢</sup> شعبة طب وجراحة العيون/ مدينة الامامين الكاظمين (ع) الطبية/ دائرة صحة بغداد/ الكرخ/ العراق.

<sup>٣</sup> وحدة الصحة العامة/ مستشفى غازي الحريري للجراحات التخصصية/ بغداد/ العراق.

<sup>٤</sup> كلية الطب/ جامعة دلتا / المنصورة الجديدة/ مصر.